## Tramadol Rapid Test Strip (Urine)

Cat.No.: TML-6081

#### INTENDED USE

The TML Rapid Test Strip (Urine) is a rapid visual immunoassay for the qualitative, presumptive detection of Tramadol in human urine specimens at the cut-off concentrations listed below:

 Parameter
 Calibrator
 Cut-off (ng/mL)

 TML (Tramadol)
 Cis-Tramadol
 100

## INTRODUCTION

Tramadol is a quasi-narcotic analgesic used in the treatment of moderate to severe pain. It is a synthetic analog of codeine, but has a low binding affinity to the mu-opioid receptors. Large doses of tramadol can develop tolerance and physiological dependency and lead to its abuse. Tramadol is extensively metabolized after oral administration. Approximately 30% of the dose is excreted in the urine as unchanged drug, whereas 60% is excreted as metabolites. The major pathways appear to be N- and O-demethylation, glucoronidation or sulfation in the liver.

#### PRINCIPLE

The TML Rapid Test Strip (Urine) detects Tramadol through visual interpretation of color development on the strip. Drug conjugates are immobilized on the test region of the membrane. During testing, the specimen reacts with antibodies conjugated to colored particles and precoated on the sample pad. The mixture then migrates through the membrane by capillary action, and interacts with reagents on the membrane. If there are insufficient drug molecules in the specimen, the antibody-colored particle conjugate will bind to the drug conjugates, forming a colored band at the test region of the membrane. Therefore, a colored band appears in the test region when the urine is negative for the drug. If drug molecules are present in the urine above the cut-off concentration of the test, they compete with the immobilized drug conjugate on the test region for limited antibody binding sites. This will prevent attachment of the antibody-colored particle conjugate to the test region. Therefore, the absence of a colored band at the test region indicates a positive result. The appearance of a colored band at the control region serves as a procedural control, indicating that the proper volume of specimen has been added and membrane wicking has occurred.

#### REAGENTS

Each test consists of a reagent strip. The amount of each antigen and/or antibody coated on the strip is less than 0.001 mg for antigen conjugates and goat anti-rabbit IgG antibodies, and less than 0.0015 mg for antibody components.

The control zone of each test contains goat anti-rabbit IgG antibody. The test zone of each test contains drug-bovine protein antigen conjugate, and the conjugate pad of each test contains monoclonal anti-drug antibody and rabbit antibody-colored particle complex

## MATERIALS

## Materials Provided

Test strips (individually pouched or in canisters)

· Package insert

### Materials Required but Not provided

· Positive and negative controls

Timer

Centrifuge

#### PRECAUTIONS

- · For professional in vitro diagnostic use only.
- Do not use after the expiration date indicated on the package. Do not use the test if the foil pouch or canister is damaged. Do not reuse tests.
- This kit contains products of animal origin. Certified knowledge of the origin and/or sanitary state
  of the animals does not completely guarantee the absence of transmissible pathogenic agents. It is
  therefore, recommended that these products be treated as potentially infectious, and handled by
  observing usual safety precautions (e.g., do not ingest or inhale).
- Avoid cross-contamination of specimens by using a new specimen collection container for each specimen obtained.
- Read the entire procedure carefully prior to testing.
- Do not eat, drink or smoke in the area where specimens and kits are handled. Handle all specimens
  as if they contain infectious agents. Observe established precautions against microbiological hazards
  throughout the procedure and follow standard procedures for the proper disposal of specimens.
  Wear protective clothing such as laboratory coats, disposable gloves and eye protection when
  specimens are assayed.
- Humidity and temperature can adversely affect results.
- Used testing materials should be discarded in accordance with local regulations.

#### STORAGE AND STABILITY

- The kit should be stored at 4-30°C until the expiry date printed on the sealed pouch.
- The test must remain in the sealed pouch or closed canister until use.
- Do not freeze.
- · Kits should be kept out of direct sunlight.
- Care should be taken to protect the components of the kit from contamination. Do not use if there is
  evidence of microbial contamination or precipitation. Biological contamination of dispensing
  equipment, containers or reagents can lead to false results.

#### SPECIMEN COLLECTION AND STORAGE

- The TML Rapid Test Strip (Urine) is intended for use with human urine specimens only.
- · Urine collected at any time of the day may be used.
- Urine specimens must be collected in clean, dry containers.
- Turbid specimens should be centrifuged, filtered, or allowed to settle and only the clear supernatant should be used for testing.
- Perform testing immediately after specimen collection. Do not leave specimens at room temperature for prolonged periods. Urine specimens may be stored at 2-8°C for up to 2 days. For long term storage, specimens should be kept below -20°C.
- Bring specimens to room temperature prior to testing. Frozen specimens must be completely thawed
  and mixed well prior to testing. Avoid repeated freezing and thawing of specimens.
- If specimens are to be shipped, pack them in compliance with all applicable regulations for transportation of etiological agents.

#### PROCEDURE

#### Bring tests, specimens, buffer and/or controls to room temperature (15-30°C) before use.

- Remove the test from its sealed pouch, or remove one strip from the canister, and use it as soon as
  possible. For best results, the assay should be performed within one hour. Canisters should be
  closed tightly after removing strips.
- Hold the strip by the end, where the product name is printed. To avoid contamination, do not touch the strip membrane.
- Holding the strip vertically, dip the test strip in the urine specimen for at least 10-15 seconds. Do not immerse past the maximum line (MAX) on the test strip.
- 4. After the test has finished running, remove the strip from the specimen and place it on a non-absorbent flat surface. Start the timer and wait for the colored band(s) to appear. The result should be read at 5 minutes. Do not interpret the result after 8 minutes.

## INTERPRETATION OF RESULTS

POSITIVE: Only one colored band appears, in the control region (C). No apparent colored band appears in the test region (T).

NEGATIVE: Two colored bands appear on the membrane. One band appears in the control region (C) and another band appears in the test region (T).



INVALID: Control band fails to appear. Results from any test which has not produced a control band at the specified read time must be discarded. Please review the procedure and repeat with a new test. If the problem persists, discontinue using the kit immediately and contact your local distributor.

#### NOTE:

- The intensity of color in the test region (T) may vary depending on the concentration of analytes present in the specimen. Therefore, any shade of color in the test region should be considered positive. Note that this is a qualitative test only, and cannot determine the concentration of analytes in the specimen.
- Insufficient specimen volume, incorrect operating procedure or expired tests are the most likely reasons for control band failure.

## QUALITY CONTROL

- Internal procedural controls are included in the test. A colored band appearing in the control region
   (C) is considered an internal positive procedural control, confirming sufficient specimen volume and correct procedural technique.
- External controls are not supplied with this kit. It is recommended that positive and negative
  controls be tested as a good laboratory practice to confirm the test procedure and to verify proper
  test performance.

## LIMITATIONS OF THE TEST

- The TML Rapid Test Strip (Urine) is for professional in vitro diagnostic use, and should be only used for the qualitative detection of Tramadol.
- 2. This assay provides a preliminary analytical test result only. A more specific alternative chemical method must be used in order to obtain a confirmed analytical result. Gas chromatography/mass spectrometry (GC/MS) has been established as the preferred confirmatory method by the National Institute on Drug Abuse (NIDA). Clinical consideration and professional judgment should be

applied to any test result, particularly when preliminary positive results are indicated.

- There is a possibility that technical or procedural errors as well as other substances and factors may interfere with the test and cause false results.
- Adulterants, such as bleach and/or alum, in urine specimens may produce erroneous results regardless of the analytical method used. Therefore, please preclude the possibility of urine adulteration prior to testing.
- A positive result indicates the presence of a Tramadol only, and does not indicate or measure intoxication.
- A negative result does not at any time rule out the presence of Tramadol in urine, as they may be present below the minimum detection level of the test.
- This test does not distinguish between Tramadol and certain medications.

#### PERFORMANCE CHARACTERISTICS

#### A. Accuracy

The accuracy of the TML Rapid Test Strip (Urine) was compared and checked against commercially available tests with a threshold value at the same cut-off levels. Urine samples taken from volunteers claiming to be non-users were examined under both tests. The results were >99.9% in agreement.

#### B. Reproducibility

The reproducibility of the TML Rapid Test Strip (Urine) was verified by blind tests performed at four different locations. Samples with Tramadol concentrations at 50% of the cut-off were all determined to be negative, while samples with Tramadol concentrations at 200% of the cut-off were all determined to be positive.

#### C. Precision

Test precision was determined by blind tests with control solutions. Controls with Tramadol concentrations at 50% of the cut-off yielded negative results, and controls with Tramadol concentrations at 150% of the cut-off yielded positive results.

#### D. Specificity

The following tables list the concentrations of compounds (ng/mL) above which the TML Rapid Test Strip (Urine) identified positive results at 5 minutes.

Tramadol related compounds	Concentration (ng/ml)
Tramadol	100
(+/-)Chlorpheniramine	50,000
Dimenhydrinate	50,000
Diphenhydramine	50,000
Phencyclidine	50,000
(+)-Chlorpheniramine	100,000

# The following compounds yielded negative results up to a concentration of 100 $\mu g/\text{mL}\colon$

(-)Cotinine	Delorazepam	Metoclopramide
(-)Isoproterenol	Desipramine HCl	Metoprolol
(+)-cis-Diltiazem	Dexamethasone	Metronidazole
(+)-Naproxen	Dextromethorphan	MOR-3-Beta-D Glucuronide
(+/-) Phenylpropanolamine	Diacetylmorphine	Nalorphine
(+/-)Epinephrine	Diazepam	Naloxone
3,4-Methylenedioxyamphetamine	Diclofenac	Nifedipine
3,4-Methylenedioxy-methamphetamine	Dicumarol	Nimesulide
4-Dimethylaminoantipyrine	Diflunisal	Nitrazepam
Acetaminophen	Digoxin	N-Methylephedrin
Acetophenetidine	Dihydrocodeine	Olanzapine
Acetylcodeine	Dimenhydrinate	Opipramol
Acetylsalicylic acid	Diphenhydramine	Oxalic acid
Alprazolam	DL-Propanolol	Oxazepam
Amikacin	DL-Tryptophan	Oxycodone
Aminopyrine	DL-Tyrosine	Oxymetazoline
Amitriptyline	Dopamine	Pennicilline G
Amoxicilline	Doxepin	Perphenazine
Amphetamine	Doxylamine	Pheniramine
Ampicilline	d-Propoxyphene	Phenothiazine
Apomorphine	Ecgonine HCl	Phentermine
Ascorbic acid	Ecgonine methylester	Prednisolone
Aspartame	Ephedrine	Prednisone
Atropine	Erythromycine	Procaine
Baclofen	Estron 3 sulfate	Promazine
Benzocaine	Ethylmorphine	Promethazine
beta-phenylethylamine	Etodolac	Prothipendyl

Bilirubin Fenfluramine Protriptyline Fentanyl Quetiapine Bromazepam Buprenorphine Fluoxetine Quinidine Caffeine Flupentixol Ranitidine Cannabidiol Furosemide Rifampicine Cannabinol Gastrozepin Risperidone Carbamazepine Gentamicin Salbutamol Chloramphenicol Gentisic acid Salicylic acid Chlordiazepoxide Guaiacol Glyceryl Ether Secobarbital Chloroquine Hemoglobin Sertraline Chlorpheniramine Hydralazine Spironolactone Chlorprothixene Hydrochlorothiazide Sulfamethoxazole Cholesterol Hydrocodone Sulindac Chorptothixene Hydrocortisone Temazepam Cimetidine Ibuprofen Thebaine Ciprofloxacin Imipramine Theophylline Citalopram Ketamine Thiamine Clindamycin Thioridazine Ketoprofen Clobazam L - Thyroxine Tobramycin Clomipramine Lidocaine Triamterene Clonazepam Lincomycin Trimethoprim Clonidine Loperamide Trimipramine Clorazepate L-Phenylephrine Tyramine Clozapine Maprotiline Vancomycin Venlafaxine Cocain Meperidine Mephentermine Codein Verapamil hemisulfate salt Creatinine Methadone Zolpidem Cyclobenzaprine Methamphetamine

## LITERATURE REFERENCES

- Baselt RC. Disposition of Toxic Drugs and Chemicals in Man. 2nd ed. Davis: Biomedical Publications; 1982.
- Hawks RL, Chiang CN, eds. Urine Testing for Drugs of Abuse. Rockville: Department of Health and Human Services, National Institute on Drug Abuse; 1986.
- Substance Abuse and Mental Health Services Administration. Mandatory Guidelines for Federal Workplace Drug Testing Programs. 53 Federal Register; 1988.
- McBay AJ. Drug-analysis technology--pitfalls and problems of drug testing. Clin Chem. 1987 Oct; 33 (11 Suppl): 33B-40B.
- Gilman AG, Goodman LS, Gilman A, eds. Goodman and Gilman's The Pharmacological Basis of Therapeutics. 6th ed. New York: Macmillan; 1980.
   GLOSSARY OF SYMBOLS

#### 

European Community

## MANUFACTURER

## ATLAS LINK TECHNOLOGY CO., LTD

Road Xing Min, Guan South Industry Zone, 065500 Langfang City, Hebei Province, CHINA WEB: https://www.invitro-test.com

CE making according to IVD Medical Directive 98/97/EC